

AQUATIC CHEMISTRY

Chemical Equilibria and Rates in Natural Waters

Third Edition

WERNER STUMM

EAWAG, Swiss Federal Institute of Technology (ETH), Zurich

JAMES J. MORGAN

Environmental Engineering Science, California Institute of Technology



A WILEY-INTERSCIENCE PUBLICATION

JOHN WILEY & SONS, INC.

New York • Chichester • Brisbane • Toronto • Singapore

SERI

Environ

We are in the third
graphs in Environ
of publishing out
ences and enginee
graduate students
has also provided
mental areas.

As new editors
and to emphasize
science. We publi
nology as it is bro
toxicology) of the
(water and wastew
ous wastes). The s
processes, the pre
remediation techn

There is a new
problems only loc
no such place as
body else in the g
sources as never
environmental deg
ecology" is emer
surroundings.

In the next 50 y
developing countr
will require a gros
will create new pr
there are new opp
for sustainable dev
erations at least
understand the en

This text is printed on acid-free paper.

Copyright © 1996 by John Wiley & Sons, Inc.

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc., 605 Third Avenue, New York, NY 10158-0012.

Library of Congress Cataloging in Publication Data:

Stumm, Werner

Aquatic chemistry: chemical equilibria and rates in natural waters / Werner Stumm, James J. Morgan.—3rd ed.

p. cm.—(Environmental science and technology)

"A Wiley-Interscience publication."

Includes bibliographical references and index.

ISBN 0-471-51184-6 (acid-free).—ISBN 0-471-51185-4 (pbk.: acid-free)

1. Water chemistry. I. Morgan, James J. II. Title.

III. Series.

GB855.S78 1995

359.9—dc20


94-48319

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

Reactions with larger ΔH^\ddagger values are affected more strongly by temperature variation. For example, for $\Delta H^\ddagger = 120 \text{ kJ mol}^{-1}$, lowering temperature from 30°C to 5°C slows the rate by a factor of 70, whereas for $\Delta H^\ddagger = 60 \text{ kJ mol}^{-1}$, the same temperature change slows the rate by a factor of about 9.

Example 2.6. Hydration of CO_2 in Terms of Transition-State Parameters A reaction of importance in natural waters and biochemistry, the hydration of CO_2 ,




is described kinetically by the parameters $E_a = 63.0$, $\Delta H^\ddagger = 60.5 \text{ (kJ mol}^{-1}\text{)}$, $\Delta S^\ddagger = -107 \text{ J K}^{-1} \text{ mol}^{-1}$, and $\log k_2 = -3.3 \text{ (M}^{-1} \text{ s}^{-1}\text{)}$ at 25°C . The slowness of this reaction ($t_{1/2} \approx 3 \text{ min}$ at 10°C) is associated with a large energy barrier as well as a highly negative activation entropy. The free energy of activation, ΔG^\ddagger , is 92 kJ mol^{-1} , giving $K^\ddagger \approx 10^{-16} \text{ M}^{-1}$, and indicating that the quasi-equilibrium concentration of the activated complex, $[\text{H}_2\text{O}, \text{CO}_2]^\ddagger$, for a 10^{-5} M CO_2 solution would be $\sim 10^{-19} \text{ M}$! For the corresponding dehydration reaction, $\text{H}_2\text{CO}_3 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$, experimental values of the *uni-molecular* rate constant vary from about 3 to 30 s^{-1} over the temperature range 5 – 30°C . Interpretation of these data in terms of ACT yields $\Delta H^\ddagger = 61.5 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -15 \text{ J K}^{-1} \text{ mol}^{-1}$ for dehydration of H_2CO_3 , consistent with $\Delta H^\circ = -1$ and $\Delta S^\circ = -92$ for the reversible *overall* reaction, and with an equilibrium constant, $K = k_f/k_b$, equal to the ratio of the hydration and dehydration rate constants, a consequence of the principle of detailed balancing (microscopic reversibility) (Moore and Pearson, 1981).

2.17. EQUILIBRIUM VERSUS STEADY STATE IN FLOW SYSTEMS

Open Flow Systems Versus Closed Systems

Most natural water systems are continuous, open systems. Flows of matter and energy occur in the real system. The time-invariant state of a continuous system with flows at the boundaries is the *steady state*.[†] This state may be poorly approximated by the *equilibrium* state of a closed system. In Figure 2.2 we indicated the important features of an open-system model with material fluxes and chemical reactions. The simple reversible reaction (a “model” reaction),



[†]The term *steady state* as used in this example means the time-invariant state of a flow system with chemical reactions. *Steady state*, with respect to chemical mechanisms, means that certain intermediates in a complex reaction are of low concentration, so that $dC/dt \approx 0$. It is important to keep these usages of “steady state” distinct.

9 $\text{Fe}^{2+}(\text{aq}) + \text{H}_2\text{O}_2 \rightarrow$	1.8	31	-116	Wells and Salam (1960)
10 $\text{Fe}^{2+} + \text{FeOH}^{2+} \rightarrow$	3.4 (21.6°C)	31	-75	Burgess (1978)
11 $\text{Fe}^{3+} + \text{Cr}^{2+} \rightarrow$	3.4	24	-117	Burgess (1978)
12 $\text{FeOH}^{2+} + \text{Cr}^{2+} \rightarrow$	6.5	22	-54	Burgess (1978)
13 $\text{Co}(\text{NH}_3)_6\text{Cl}^{2+} + \text{Fe}^{2+} \rightarrow$	-2.9	55	-126	Burgess (1978)
Dissociation:				
14 $\text{H}_2\text{CO}_3 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$	1.2 s ⁻¹	64	-15	Edsall (1969)

^a25°C, except as noted.

^bM⁻¹ s⁻¹, except for reaction 14, unimolecular, s⁻¹.

^cSolvent exchange formulated as second order, for first order, $k = k_{\text{SS}}/6$, that is, $[\text{H}_2\text{O}] = 55 \text{ M}$.

was introduced to illustrate elementary differences between closed- and open-system models. Solution for the steady-state concentration values of A and B gives the results

$$C_A = \frac{r\bar{C}_{A,0} + k_b(\bar{C}_{A,0} + \bar{C}_{B,0})}{k_f + k_b + r} \quad (13)$$

$$C_B = \frac{r\bar{C}_{B,0} + k_f(\bar{C}_{A,0} + \bar{C}_{B,0})}{k_f + k_b + r} \quad (14)$$

in which $r = Q/V$, the fluid flow rate constant (time^{-1}), and overbars denote inflowing concentrations.

We examine the ratio of C_B and C_A and compare it with the ratio expected for chemical equilibrium. The ratio of C_B/C_A for the steady state is, dividing equation 139 by equation 138,

$$\frac{C_B}{C_A} = \frac{r\bar{C}_{B,0} + k_f(\bar{C}_{A,0} + \bar{C}_{B,0})}{r\bar{C}_{A,0} + k_b(\bar{C}_{A,0} + \bar{C}_{B,0})} \quad (14)$$

Equation (140) shows that C_B/C_A will tend toward k_f/k_b as the material flow rate to the system becomes small, that is, as $r\bar{C}_{B,0}$ and $r\bar{C}_{A,0}$ vanish. For $r = 0$, the system becomes a closed system, and $C_B/C_A = k_f/k_b = K$, the equilibrium constant. The quantity $r = Q/V$ is the reciprocal of the fluid residence time of the well-mixed system: $r = \tau_R^{-1}$. As τ_R tends to very large values, the steady-state concentrations of the system approach the equilibrium values.

A simple result is obtained when only A enters the system. Then, $\bar{C}_{B,0} = 0$ and equation 140 reduces to

$$\frac{C_B}{C_A} = \frac{k_f\bar{C}_{A,0}}{r\bar{C}_{A,0} + k_b\bar{C}_{A,0}} = \frac{k_f}{r + k_b}$$

The steady-state concentration ratio quotient depends on the chemical rate constants and the flux rate constant. For $r \ll k_b$, $C_B/C_A \approx k_f/k_b = K$; if $r \gg k_b$, then $C_B/C_A \approx k_f/r$. In terms of the residence time, τ_R , and the half-time of the backward reaction, τ_b , the steady-state ratio approximates the equilibrium ratio if $\tau_R \gg \tau_b$.

Example 2.7. Steady-State Composition of an Open Completely Mixed System with the Reaction $A \rightleftharpoons B$ as a Function of Residence Time Assume $\bar{C}_{B,0} = 0$, $k_f = 10^{-5} \text{ s}^{-1}$, and $k_b = 10^{-6} \text{ s}^{-1}$, so that $K = 10$. The residence time, τ_R , is varied over the range from 10^5 to 10^8 s . Table 2.10 shows the calculated ratio of steady-state C_B/C_A to K for a 1000-fold variation in residence time relative to the characteristic reaction time ($\sim k_b^{-1}$).

The equilibrium assumption is justified for many reactions with short τ_R and long τ_b . Rate data for a large number of first- and second-order aqueous

Table 2.9. Kinetic Parameters for Elementary Aqueous Reactions^a

Reactants	$\log k^0$ ($\text{M}^{-1} \text{ s}^{-1}$)	E_a (kJ mol^{-1})	ΔS^\ddagger ($\text{J K}^{-1} \text{ mol}^{-1}$)	References
-----------	--	-----------------------------------	---	------------

reactions indicate reactions have times of a number and Lerman, 19 reactions for which the time-in of local equilibrium of local equilibrium condition species and phase environments. In (hetically active biologically active total system equilibrium

SUGGESTED

Atkins, P. W. (19 Berner, R. A. (197 Blandamer, M. J. Denbigh, K. G. (1 Chemistry and bridge. Edsall, J. T. (196 Properties and Physiology DC.

Table 2.10. Composition^a of a Completely Mixed Open System at Steady State for $A \xrightleftharpoons[k_b]{k_f} B$

τ_R (s)	r (s^{-1})	r/k_b	C_B/C_A K
10^5	10^{-5}	10	0.09
2×10^5	5×10^{-6}	5	0.17
5×10^5	2×10^{-6}	2	0.33
10^6	1×10^{-6}	1	0.50
2×10^6	5×10^{-7}	0.5	0.67
5×10^6	2×10^{-7}	0.2	0.83
10^7	1×10^{-7}	0.1	0.91
10^8	1×10^{-8}	0.01	0.99

^a $(C_B/C_A)/K$ versus τ_R ; $k_f = 1 \times 10^{-5} s^{-1}$; $k_b = 1 \times 10^{-6} s^{-1}$.

reactions indicate τ_{chem} less than seconds to minutes, and many other aqueous reactions have τ_{chem} less than hours to days (Hoffmann, 1981). The residence times of a number of freshwater systems are greater than these ranges (Imboden and Lerman, 1979). Critical attention needs to be directed to slow chemical reactions for which $\tau_{chem} \geq \tau_R$.

For many systems it is known that there exist regions or environments in which the time-invariant condition closely approaches equilibrium. The concept of local equilibrium is important in examining complex systems. Local equilibrium conditions are expected to develop, for example, for kinetically rapid species and phases at sediment-water interfaces in fresh, estuarine, and marine environments. In contrast, other local environments, such as the photosynthetically active surface regions of nearly all lakes and ocean waters and the biologically active regions of soil-water systems, are clearly far removed from total system equilibrium.

SUGGESTED READINGS

- Atkins, P. W. (1990) *Physical Chemistry*, 4th ed., Freeman, New York.
- Berner, R. A. (1971) *Principles of Chemical Sedimentology*, McGraw-Hill, New York.
- Blandamer, M. J. (1992) *Chemical Equilibria in Solution*, Ellis Horward, New York.
- Denbigh, K. G. (1971) *The Principles of Chemical Equilibrium: With Applications in Chemistry and Chemical Engineering*, 3rd ed., Cambridge University Press, Cambridge.
- Edsall, J. T. (1969) Carbon Dioxide, Carbonic Acid and Bicarbonate Ion: Physical Properties and Kinetics of Interconversion. In *CO₂: Chemical, Biochemical and Physiological Aspects*, Forster, R. E., et al., Eds., NASA SP-188, Washington, DC.